


(Estd. 1950)

Science College, Nanded

Reaccredited with 'A' Grade by NAAC with CGPA 3.38

College with Potential for Excellence (CPE Status) / DST-FIST Programme

Best College Award by S. R. T. M. U., Nanded



National Conference on "EMERGING TRENDS IN CHEMICAL SCIENCES" NCETCS-2019

20th - 21st September 2019

SOUVENIR

Organised By

Department of Chemistry, Analytical Chemistry,
Agrochemical and Fertilizers.

N.E.S. Science College

Nanded 431605, Maharashtra India | Visit us www.sciencecollegennd.org.

150

UGC-CPE & S.R.T.M. University, Nanded.

Page No	Titles and Authors Name
68	Phytochemical and HPTLC Studies on <i>Tridax procumbens</i> -A common medicinal plant of Asteriaceae Sandip Padwal and D. M. Jadhav
69	Synthesis and Characterization of new Transition Metal Complexes Derived from Schiff Bases Lande A.A.*¹, Kolapwar B.G.*¹, Karamunge K.G.*²
70	Microwave assisted synthesis and antimicrobial study of some novel 2-azetidinones derived from 2-(1-Phenylimino-ethyl)-naphthalen-1-ol Yogesh Nalwar¹, Sainath Zangade², Avinash Shinde³ and Pravinkumar Patil
71	Investigation of volumetric and acoustic properties of Promethazine hydrochloride in aqueous L-Proline solution at different temperature R. T. Sawale¹, A. D. Arsule², A. V. Pawde³ and S. D. Deosarkar
72	Validation of reverse phase hplc method for the estimation of metformin and glibenclamide in tablet dosage form Wayal Shubhangi¹, T. M. Kalyankar² and Anitha Kowthalam
73	Cyanuric chloride catalyzed three component, one pot synthesis of Biginelli-type pyrimidinone derivatives Baliram S. Hote*, Gajanan G. Mandawad, Sudhakar G. Patil, Shivshankar N. Hallale
74	Lanthanum (III) nitrate catalyzed one pot synthesis of dipyrromethanes at room temperature Kabeer A. Shaikh and Uddhav N. Chaudhar
RESEARCH ARTICLES	
75	Microwave Assisted Facile and Efficient One-Pot Synthesis of 2,3-Dihydroquinazolin-4(1H)-Ones in Ionic Liquid [bmim]HSO ₄ Under Solvent Free Conditions. Sandeep V. Shinde, Rajesh H. Tale, Arjun V. Nakhate
89	Synthesis, Characterization and Biological Potential of Some New 4-Fluorochalcones in Water under Ultrasonication Balasaheb B. Musle, Satish V. Gaikwad, Nana V. Shitole, Suraj B. Ade and Subhash M. Lonkar
95	Determination of saponification value of oils by microwave irradiation technique Suresh D. Dhage, D.R. Munde and Swanand Mukhedkar

Determination of saponification value of oils by microwave irradiation technique

Suresh D. Dhage¹; D.R. Munde² and Swanand Mukhedkar³

¹Dept. of Chemistry, SSJES, Arts, Commerce and Science College, Gangakhed-431514.

²Dept. of Chemistry, NES's, Science College, Nanded

³Dept. of Chemistry, Shahir Annabhau Sathe College, Mukhed. Dist. Nanded

E-mail: dhage137@gmail.com

ABSTRACT

A rapid, simple and accurate method for the determination of saponification values of fixed oils are reported under microwave irradiation. The saponification reaction was completed within 5 min. and similar results were obtained, when compared with conventional methods.

Key words: Microwave energy, Fixed oils, Saponification value

INTRODUCTION

In recent times, the use of microwave irradiation is a crucial matter in our scientific community. Most of the industrial as well as academic research groups are utilizing the techniques of microwave-assisted organic synthesis as the latest technology for rapid optimization of reaction, for the efficient synthesis of new chemical entities and also for discovering new chemical reactivity. It is a useful tool towards Green chemistry, which helps in minimizing environmental pollution. Microwave induced Organic Reaction Enhancement chemistry (MORE) is gaining popularity as a nonconventional technique for rapid organic synthesis. It can be termed as 'e-chemistry' because it is easy, effective, economic and ecofriendly and is believed to be a step towards Green chemistry¹. Short response time and highly accelerated reaction rate are main advantages of MORE chemistry². Microwave heating reduces side reactions, increases percentage yield and it improves reproducibility^{3,4}. In the proposed method, the microwave technique has been applied for the determination of saponification value of several fixed oils. The conventional pharmacopoeias procedure involves thirty minutes heating of the reaction mixture^{5,6}. Through this, we have developed a simple accurate and rapid procedure for the determination of saponification value.

EXPERIMENTAL

Fresh oils were obtained from the refineries of known source. A mixture of 2 g of oil sample and 25 mL of 0.5M alcoholic KOH was taken in an open Borosil beaker covered with a Petri dish containing few ice pieces (to avoid excessive evaporation of solvent). It was placed in a domestic microwave oven and subjected for irradiation at 60 watts microwave intensity for 5 min. A 500 mL

beaker containing 200 mL water was placed in the oven next to the reaction beaker to serve as heating sink⁷. The reaction mixture was cooled and titrated against 0.5 M hydrochloric acid using phenolphthalein indicator. A blank was also carried out. The saponification value of each sample was determined using the formula given in the pharmacopoeia. The procedure was repeated thrice for each sample and mean values were compared with conventional and standard values⁸⁻¹⁰(Table 1).

Table 1: Determination of saponification value

Sample	Standard Values	Conventional method	Microwave method
Cottonseed oil	190-198	192.2	192.9
Olive oil	182-195	189.7	187.4
Almond oil	181-203	201.4	202.7
Palm oil	187-201	191.8	193.5
Sesame oil	185-194	186.2	192.6
Safflower oil	186-197	191.4	193.9
Groundnut oil	188-195	189.9	194.1
Corn oil	187-194	186.3	191.4
Codliver oil	181-189	187.6	183.5

RESULTS AND DISCUSSION

Under microwave conditions, the reaction is rapid (5 minutes) compared to the conventional method (30 minutes). The saponification values for all the samples tested were similar to their conventional and standard values. The developed method is a rapid, simple, accurate and efficient method for the determination of a large number of oil samples, waxes and surfactants. It can also be used in industries and academic institutions for the routine analysis and quality control of oils and waxes. It is expected that in this century, most of the chemists will probably use microwave energy to heat chemical reaction on a laboratory scale.

REFERENCES

1. S. V. Sharma, G. V. S. Ramasarma and B. Suresh, *Ind. J. Pharm. Sci.*, **64**, 337-344 (2002).
2. R. N. Gedey, F. E. Smith, K. C. Westaway, H. Ali, L. Baldisera, L. Laberge and J. Rousell, *Tetrahedron Lett.*, **27**, 279 (1986).

3. Subarna Ganguly, *Pharma Times*, 38 (12) , 26, (2006).
4. G. Richard, S. Frank, W. Kenneth, A. Humera, B. Lorraine and L. Lena, *Tetrahedron Lett.*, 27, 279 (1986).
5. A. K. Bose, M. S. Manhas, M. Ghosh, M. Shah, V. S. Raju, S. S. Bari, S. N. Newaz, B. K. Banic, A. G. Choudhari and K. J. Barkat, *J. Org. Chem.*, 56, 6968 (1991).
6. *The Pharmacopoeia of India*, Govt. of India, Ministry of Health, The Manager of Publication, New Delhi, (1996) p. 936.
7. S. Badami, S. A. Manohara Reddy, S. V. Sharma and B. Suresh, *Indian Drugs*, 39, 451-452 (2002).
8. W. C. Evans, *Textbook of Pharmacognosy*, Saunders Company Limited, Edinburgh, (1996) p.181.
9. Ronald S. Kirk and Ronald Sawyer, *Pearson's Composition and Analysis of Foods*, Addison Wesley Longman Ltd., (1996) p. 617.
10. T. E. Wallis, *Textbook of Pharmacognosy*, C B S Publishers and Distributors, New Delhi, (1997) p. 210.